

Frontiers in Pharmacology, 2018, vol.9, NMAR

Application of Mesenchymal stem cells for therapeutic agent delivery in anti-tumor treatment

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Chulpanova, Kitaeva, Tazetdinova, James, Rizvanov and Solovyeva. Mesenchymal stem cells (MSCs) are non-hematopoietic progenitor cells, which can be isolated from different types of tissues including bone marrow, adipose tissue, tooth pulp, and placenta/umbilical cord blood. Their isolation from adult tissues circumvents the ethical concerns of working with embryonic or fetal stem cells, whilst still providing cells capable of differentiating into various cell lineages, such as adipocytes, osteocytes and chondrocytes. An important feature of MSCs is the low immunogenicity due to the lack of co-stimulatory molecules expression, meaning there is no need for immunosuppression during allogeneic transplantation. The tropism of MSCs to damaged tissues and tumor sites makes them a promising vector for therapeutic agent delivery to tumors and metastatic niches. MSCs can be genetically modified by virus vectors to encode tumor suppressor genes, immunomodulating cytokines and their combinations, other therapeutic approaches include MSCs priming/loading with chemotherapeutic drugs or nanoparticles. MSCs derived membrane microvesicles (MVs), which play an important role in intercellular communication, are also considered as a new therapeutic agent and drug delivery vector. Recruited by the tumor, MSCs can exhibit both pro- and anti-oncogenic properties. In this regard, for the development of new methods for cancer therapy using MSCs, a deeper understanding of the molecular and cellular interactions between MSCs and the tumor microenvironment is necessary. In this review, we discuss MSC and tumor interaction mechanisms and review the new therapeutic strategies using MSCs and MSCs derived MVs for cancer treatment.

<http://dx.doi.org/10.3389/fphar.2018.00259>

Keywords

Chemotherapy resistance, Cytokines, Membrane vesicles, Mesenchymal stem cells, Oncolytic viruses, Suppressor genes, Tumor microenvironment

References

- [1] Ahn, J., Lee, H., Seo, K., Kang, S., Ra, J., and Youn, H. (2013). Anti-tumor effect of adipose tissue derived-mesenchymal stem cells expressing interferon-beta and treatment with cisplatin in a xenograft mouse model for canine melanoma. *PLoS One* 8:e74897. doi: 10.1371/journal.pone.0074897
- [2] Arvelo, F., Sojo, F., and Cotte, C. (2016). Tumour progression and metastasis. *Ecancermedicalscience* 10:617. doi: 10.3332/ecancer.2016.617
- [3] Birnbaum, T., Roeder, J., Schankin, C. J., Padovan, C. S., Schichor, C., Goldbrunner, R., et al. (2007). Malignant gliomas actively recruit bone marrow stromal cells by secreting angiogenic cytokines. *J. Neurooncol.* 83, 241-247. doi: 10.1007/s11060-007-9332-4

- [4] Blatt, N. L., Mingaleeva, R. N., Khaiboullina, S. F., Kotlyar, A., Lombardi, V. C., and Rizvanov, A. A. (2013a). In vivo screening models of anticancer drugs. *Life Sci. J.* 10, 1892-1900.
- [5] Blatt, N. L., Mingaleeva, R. N., Khaiboullina, S. F., Lombardi, V. C., and Rizvanov, A. A. (2013b). Application of cell and tissue culture systems for anticancer drug screening. *World Appl. Sci. J.* 23, 315-325. doi: 10.5829/idosi.wasj.2013.23.03.13064
- [6] Bliss, S. A., Sinha, G., Sandiford, O. A., Williams, L. M., Engelberth, D. J., Guiro, K., et al. (2016). Mesenchymal stem cell-derived exosomes stimulate cycling quiescence and early breast cancer dormancy in bone marrow. *Cancer Res.* 76, 5832-5844. doi: 10.1158/0008-5472.CAN-16-1092
- [7] Bolontrade, M. F., Sganga, L., Piaggio, E., Viale, D. L., Sorrentino, M. A., Robinson, A., et al. (2012). A specific subpopulation of mesenchymal stromal cell carriers overrides melanoma resistance to an oncolytic adenovirus. *Stem Cells Dev.* 21, 2689-2702. doi: 10.1089/scd.2011.0643
- [8] Bonomi, A., Ghezzi, E., Pascucci, L., Aralla, M., Ceserani, V., Pettinari, L., et al. (2017a). Effect of canine mesenchymal stromal cells loaded with paclitaxel on growth of canine glioma and human glioblastoma cell lines. *Vet. J.* 223, 41-47. doi: 10.1016/j.tvjl.2017.05.005
- [9] Bonomi, A., Steimberg, N., Benetti, A., Berenzi, A., Alessandri, G., Pascucci, L., et al. (2017b). Paclitaxel-releasing mesenchymal stromal cells inhibit the growth of multiple myeloma cells in a dynamic 3D culture system. *Hematol. Oncol.* 35, 693-702. doi: 10.1002/hon.2306
- [10] Brini, A. T., Cocce, V., Ferreira, L. M., Giannasi, C., Cossellu, G., Gianni, A. B., et al. (2016). Cell-mediated drug delivery by gingival interdental papilla mesenchymal stromal cells (GinPa-MSCs) loaded with paclitaxel. *Expert Opin. Drug Deliv.* 13, 789-798. doi: 10.1517/17425247.2016.1167037
- [11] Cafforio, P., Viggiano, L., Mannavola, F., Pelle, E., Caporusso, C., Maiorano, E., et al. (2017). pIL6-TRAIL-engineered umbilical cord mesenchymal/stromal stem cells are highly cytotoxic for myeloma cells both in vitro and in vivo. *Stem Cell Res. Ther.* 8:206. doi: 10.1186/s13287-017-0655-6
- [12] Cai, C., Hou, L., Zhang, J., Zhao, D., Wang, Z., Hu, H., et al. (2017). The inhibitory effect of mesenchymal stem cells with rAd-NK4 on liver cancer. *Appl. Biochem. Biotechnol.* 183, 444-459. doi: 10.1007/s12010-017-2456-x
- [13] Chanda, D., Isayeva, T., Kumar, S., Hensel, J. A., Sawant, A., Ramaswamy, G., et al. (2009). Therapeutic potential of adult bone marrow-derived mesenchymal stem cells in prostate cancer bone metastasis. *Clin. Cancer Res.* 15, 7175-7185. doi: 10.1158/1078-0432.CCR-09-1938
- [14] Chen, Q., Cheng, P., Yin, T., He, H., Yang, L., Wei, Y., et al. (2012). Therapeutic potential of bone marrow-derived mesenchymal stem cells producing pigment epithelium-derived factor in lung carcinoma. *Int. J. Mol. Med.* 30, 527-534. doi: 10.3892/ijmm.2012.1015
- [15] Chen, X., Lin, X., Zhao, J., Shi, W., Zhang, H., Wang, Y., et al. (2008). A tumor-selective biotherapy with prolonged impact on established metastases based on cytokine gene-engineered MSCs. *Mol. Ther.* 16, 749-756. doi: 10.1038/mt.2008.3
- [16] Choi, S. A., Hwang, S. K., Wang, K. C., Cho, B. K., Phi, J. H., Lee, J. Y., et al. (2011). Therapeutic efficacy and safety of TRAIL-producing human adipose tissue-derived mesenchymal stem cells against experimental brainstem glioma. *Neuro Oncol.* 13, 61-69. doi: 10.1093/neuonc/noq147
- [17] Cocce, V., Balducci, L., Falchetti, M. L., Pascucci, L., Ciusani, E., Brini, A. T., et al. (2017a). Fluorescent immortalized human adipose derived stromal cells (hASCs-TS/GFP+) for studying cell drug delivery mediated by microvesicles. *Anticancer Agents Med. Chem.* 17, 1578-1585. doi: 10.2174/1871520617666170327113932
- [18] Cocce, V., Farronato, D., Brini, A. T., Masia, C., Gianni, A. B., Piovani, G., et al. (2017b). Drug loaded gingival mesenchymal stromal cells (GinPa-MSCs) inhibit in vitro proliferation of oral squamous cell carcinoma. *Sci. Rep.* 7:9376. doi: 10.1038/s41598-017-09175-4
- [19] Cousin, B., Ravet, E., Poglio, S., De Toni, F., Bertuzzi, M., Lulka, H., et al. (2009). Adult stromal cells derived from human adipose tissue provoke pancreatic cancer cell death both in vitro and in vivo. *PLoS One* 4:e6278. doi: 10.1371/journal.pone.0006278
- [20] Del Fattore, A., Luciano, R., Saracino, R., Battafarano, G., Rizzo, C., Pascucci, L., et al. (2015). Differential effects of extracellular vesicles secreted by mesenchymal stem cells from different sources on glioblastoma cells. *Expert Opin. Biol. Ther.* 15, 495-504. doi: 10.1517/14712598.2015.997706
- [21] Doi, C., Maurya, D. K., Pyle, M. M., Troyer, D., and Tamura, M. (2010). Cytotherapy with naive rat umbilical cord matrix stem cells significantly attenuates growth of murine pancreatic cancer cells and increases survival in syngeneic mice. *Cytotherapy* 12, 408-417. doi: 10.3109/14653240903548194
- [22] Du, J., Zhang, Y., Xu, C., and Xu, X. (2015). Apoptin-modified human mesenchymal stem cells inhibit growth of lung carcinoma in nude mice. *Mol. Med. Rep.* 12, 1023-1029. doi: 10.3892/mmr.2015.3501
- [23] Du, W., Seah, I., Bougazzoul, O., Choi, G., Meeth, K., Bosenberg, M. W., et al. (2017). Stem cell-released oncolytic herpes simplex virus has therapeutic efficacy in brain metastatic melanomas. *Proc. Natl. Acad. Sci. U.S.A.* 114, E6157-E6165. doi: 10.1073/pnas.1700363114
- [24] Duebgen, M., Martinez-Quintanilla, J., Tamura, K., Hingtgen, S., Redjal, N., Wakimoto, H., et al. (2014). Stem cells loaded with multimechanistic oncolytic herpes simplex virus variants for brain tumor therapy. *J. Natl. Cancer Inst.* 106:dju090. doi: 10.1093/jnci/dju090

- [25] Dvorak, H. F. (1986). Tumors: wounds that do not heal. Similarities between tumor stroma generation and wound healing. *N. Engl. J. Med.* 315, 1650-1659. doi: 10.1056/NEJM198612253152606
- [26] Gilazieva, Z. E., Tazetdinova, L. G., Arkhipova, S. S., Solovyeva, V. V., and Rizvanov, A. A. (2016). Effect of cisplatin on ultrastructure and viability of adipose-derived mesenchymal stem cells. *BioNanoScience* 6, 534-539. doi: 10.1007/s12668-016-0283-0
- [27] Gu, H., Ji, R., Zhang, X., Wang, M., Zhu, W., Qian, H., et al. (2016). Exosomes derived from human mesenchymal stem cells promote gastric cancer cell growth and migration via the activation of the Akt pathway. *Mol. Med. Rep.* 14, 3452-3458. doi: 10.3892/mmr.2016.5625
- [28] Guiho, R., Biteau, K., Grisendi, G., Taurelle, J., Chatelais, M., Gantier, M., et al. (2016). TRAIL delivered by mesenchymal stromal/stem cells counteracts tumor development in orthotopic Ewing sarcoma models. *Int. J. Cancer* 139, 2802-2811. doi: 10.1002/ijc.30402
- [29] Guo, X. R., Hu, Q. Y., Yuan, Y. H., Tang, X. J., Yang, Z. S., Zou, D. D., et al. (2016). PTEN-mRNA engineered mesenchymal stem cell-mediated cytotoxic effects on U251 glioma cells. *Oncol. Lett.* 11, 2733-2740. doi: 10.3892/ol.2016.4297
- [30] Hakkarainen, T., Sarkioja, M., Lehenkari, P., Miettinen, S., Ylikomi, T., Suuronen, R., et al. (2007). Human mesenchymal stem cells lack tumor tropism but enhance the antitumor activity of oncolytic adenoviruses in orthotopic lung and breast tumors. *Hum. Gene Ther.* 18, 627-641. doi: 10.1089/hum.2007.034
- [31] Han, J., Zhao, J., Xu, J., and Wen, Y. (2014). Mesenchymal stem cells genetically modified by lentivirus-mediated interleukin-12 inhibit malignant ascites in mice. *Exp. Ther. Med.* 8, 1330-1334. doi: 10.3892/etm.2014.1918
- [32] Herberts, C. A., Kwa, M. S., and Hermesen, H. P. (2011). Risk factors in the development of stem cell therapy. *J. Transl. Med.* 9:29. doi: 10.1186/1479-5876-9-29
- [33] Hong, X., Miller, C., Savant-Bhonsale, S., and Kalkanis, S. N. (2009). Antitumor treatment using interleukin-12-secreting marrow stromal cells in an invasive glioma model. *Neurosurgery* 64, 1139-1146; discussion 1146-1147. doi: 10.1227/01.NEU.0000345646.85472.EA
- [34] Hu, W., Wang, J., He, X., Zhang, H., Yu, F., Jiang, L., et al. (2011). Human umbilical blood mononuclear cell-derived mesenchymal stem cells serve as interleukin-21 gene delivery vehicles for epithelial ovarian cancer therapy in nude mice. *Biotechnol. Appl. Biochem.* 58, 397-404. doi: 10.1002/bab.63
- [35] Huang, W. H., Chang, M. C., Tsai, K. S., Hung, M. C., Chen, H. L., and Hung, S. C. (2013). Mesenchymal stem cells promote growth and angiogenesis of tumors in mice. *Oncogene* 32, 4343-4354. doi: 10.1038/onc.2012.458
- [36] Ji, R., Zhang, B., Zhang, X., Xue, J., Yuan, X., Yan, Y., et al. (2015). Exosomes derived from human mesenchymal stem cells confer drug resistance in gastric cancer. *Cell Cycle* 14, 2473-2483. doi: 10.1080/15384101.2015.1005530
- [37] Kalimuthu, S., Gangadaran, P., Li, X. J., Oh, J. M., Lee, H. W., Jeong, S. Y., et al. (2016). In Vivo therapeutic potential of mesenchymal stem cell-derived extracellular vesicles with optical imaging reporter in tumor mice model. *Sci. Rep.* 6:30418. doi: 10.1038/srep30418
- [38] Kalimuthu, S., Oh, J. M., Gangadaran, P., Zhu, L., Lee, H. W., Rajendran, R. L., et al. (2017). In vivo tracking of chemokine receptor CXCR4-engineered mesenchymal stem cell migration by optical molecular imaging. *Stem Cells Int.* 2017:8085637. doi: 10.1155/2017/8085637
- [39] Kanehira, M., Xin, H., Hoshino, K., Maemondo, M., Mizuguchi, H., Hayakawa, T., et al. (2007). Targeted delivery of NK4 to multiple lung tumors by bone marrow-derived mesenchymal stem cells. *Cancer Gene Ther.* 14, 894-903. doi: 10.1038/sj.cgt.7701079
- [40] Khakoo, A. Y., Pati, S., Anderson, S. A., Reid, W., Elshal, M. F., and Rovira, I. I., et al. (2006). Human mesenchymal stem cells exert potent antitumorigenic effects in a model of Kaposi's sarcoma. *J. Exp. Med.* 203, 1235-1247. doi: 10.1084/jem.20051921
- [41] Khorashadizadeh, M., Soleimani, M., Khanahmad, H., Fallah, A., Naderi, M., and Khorramizadeh, M. (2015). Bypassing the need for pre-sensitization of cancer cells for anticancer TRAIL therapy with secretion of novel cell penetrable form of Smac from hA-MSCs as cellular delivery vehicle. *Tumour Biol.* 36, 4213-4221. doi: 10.1007/s13277-015-3058-2
- [42] Kidd, S., Caldwell, L., Dietrich, M., Samudio, I., Spaeth, E. L., Watson, K., et al. (2010). Mesenchymal stromal cells alone or expressing interferon-beta suppress pancreatic tumors in vivo, an effect countered by anti-inflammatory treatment. *Cytotherapy* 12, 615-625. doi: 10.3109/14653241003631815
- [43] Kim, N., Nam, Y. S., Im, K. I., Lim, J. Y., Lee, E. S., Jeon, Y. W., et al. (2015). IL-21-expressing mesenchymal stem cells prevent lethal B-cell lymphoma through efficient delivery of IL-21, which redirects the immune system to target the tumor. *Stem Cells Dev.* 24, 2808-2821. doi: 10.1089/scd.2015.0103
- [44] Kim, S. M., Oh, J. H., Park, S. A., Ryu, C. H., Lim, J. Y., Kim, D. S., et al. (2010). Irradiation enhances the tumor tropism and therapeutic potential of tumor necrosis factor-related apoptosis-inducing ligand-secreting human umbilical cord blood-derived mesenchymal stem cells in glioma therapy. *Stem Cells* 28, 2217-2228. doi: 10.1002/stem.543

- [45] Klopp, A. H., Spaeth, E. L., Dembinski, J. L., Woodward, W. A., Munshi, A., Meyn, R. E., et al. (2007). Tumor irradiation increases the recruitment of circulating mesenchymal stem cells into the tumor microenvironment. *Cancer Res.* 67, 11687-11695. doi: 10.1158/0008-5472.CAN-07-1406
- [46] Kucerova, L., Altanerova, V., Matuskova, M., Tyciakova, S., and Altaner, C. (2007). Adipose tissue-derived human mesenchymal stem cells mediated prodrug cancer gene therapy. *Cancer Res.* 67, 6304-6313. doi: 10.1158/0008-5472.CAN-06-4024
- [47] Kucerova, L., Matuskova, M., Pastorakova, A., Tyciakova, S., Jakubikova, J., Bohovic, R., et al. (2008). Cytosine deaminase expressing human mesenchymal stem cells mediated tumour regression in melanoma bearing mice. *J. Gene Med.* 10, 1071-1082. doi: 10.1002/jgm.1239
- [48] Lathrop, M. J., Sage, E. K., Macura, S. L., Brooks, E. M., Cruz, F., Bonenfant, N. R., et al. (2015). Antitumor effects of TRAIL-expressing mesenchymal stromal cells in a mouse xenograft model of human mesothelioma. *Cancer Gene Ther.* 22, 44-54. doi: 10.1038/cgt.2014.68
- [49] Lee, H. K., Finniss, S., Cazacu, S., Bucris, E., Ziv-Av, A., Xiang, C., et al. (2013). Mesenchymal stem cells deliver synthetic microRNA mimics to glioma cells and glioma stem cells and inhibit their cell migration and self-renewal. *Oncotarget* 4, 346-361. doi: 10.18632/oncotarget.868
- [50] Lejmi, E., Perriraz, N., Clement, S., Morel, P., Baertschiger, R., Christofilopoulos, P., et al. (2015). Inflammatory chemokines MIP-1delta and MIP-3alpha are involved in the migration of multipotent mesenchymal stromal cells induced by hepatoma cells. *Stem Cells Dev.* 24, 1223-1235. doi: 10.1089/scd.2014.0176
- [51] Li, L., Guan, Y., Liu, H., Hao, N., Liu, T., Meng, X., et al. (2011). Silica nanorattle-doxorubicin-anchored mesenchymal stem cells for tumor-tropic therapy. *ACS Nano* 5, 7462-7470. doi: 10.1021/nn202399w
- [52] Li, L., Li, F., Tian, H., Yue, W., Li, S., and Chen, G. (2014). Human mesenchymal stem cells with adenovirus-mediated TRAIL gene transduction have antitumor effects on esophageal cancer cell line Eca-109. *Acta Biochim. Biophys. Sin.* 46, 471-476. doi: 10.1093/abbs/gmu024
- [53] Li, X., Lu, Y., Huang, W., Xu, H., Chen, X., Geng, Q., et al. (2006). In vitro effect of adenovirus-mediated human Gamma Interferon gene transfer into human mesenchymal stem cells for chronic myelogenous leukemia. *Hematol. Oncol.* 24, 151-158. doi: 10.1002/hon.779
- [54] Lin, R., Wang, S., and Zhao, R. C. (2013). Exosomes from human adipose-derived mesenchymal stem cells promote migration through Wnt signaling pathway in a breast cancer cell model. *Mol. Cell. Biochem.* 383, 13-20. doi: 10.1007/s11010-013-1746-z
- [55] Ling, X., Marini, F., Konopleva, M., Schober, W., Shi, Y., Burks, J., et al. (2010). Mesenchymal stem cells overexpressing IFN-beta inhibit breast cancer growth and metastases through Stat3 signaling in a syngeneic tumor model. *Cancer Microenviron.* 3, 83-95. doi: 10.1007/s12307-010-0041-8
- [56] Liu, S., Ginestier, C., Ou, S. J., Clouthier, S. G., Patel, S. H., Monville, F., et al. (2011). Breast cancer stem cells are regulated by mesenchymal stem cells through cytokine networks. *Cancer Res.* 71, 614-624. doi: 10.1158/0008-5472.CAN-10-0538
- [57] Ma, F., Chen, D., Chen, F., Chi, Y., Han, Z., Feng, X., et al. (2015). Human umbilical cord mesenchymal stem cells promote breast cancer metastasis by interleukin-8- and interleukin-6-dependent induction of CD44+/CD24- cells. *Cell Transplant.* 24, 2585-2599. doi: 10.3727/096368915X687462
- [58] Marini, I., Siegemund, M., Hutt, M., Kontermann, R. E., and Pfizenmaier, K. (2017). Antitumor activity of a mesenchymal stem cell line stably secreting a tumor-targeted TNF-related apoptosis-inducing ligand fusion protein. *Front. Immunol.* 8:536. doi: 10.3389/fimmu.2017.00536
- [59] Martinez-Quintanilla, J., Bhare, D., Heidari, P., He, D., Mahmood, U., and Shah, K. (2013). Therapeutic efficacy and fate of bimodal engineered stem cells in malignant brain tumors. *Stem Cells* 31, 1706-1714. doi: 10.1002/stem.1355
- [60] Matuskova, M., Hlubinova, K., Pastorakova, A., Hunakova, L., Altanerova, V., Altaner, C., et al. (2010). HSV-tk expressing mesenchymal stem cells exert bystander effect on human glioblastoma cells. *Cancer Lett.* 290, 58-67. doi: 10.1016/j.canlet.2009.08.028
- [61] Mohammadpour, H., Pourfathollah, A. A., Nikougoftar Zarif, M., and Shahbazfar, A. A. (2016). Irradiation enhances susceptibility of tumor cells to the antitumor effects of TNF-alpha activated adipose derived mesenchymal stem cells in breast cancer model. *Sci. Rep.* 6:28433. doi: 10.1038/srep28433
- [62] Mohr, A., Lyons, M., Deedigan, L., Harte, T., Shaw, G., Howard, L., et al. (2008). Mesenchymal stem cells expressing TRAIL lead to tumour growth inhibition in an experimental lung cancer model. *J. Cell Mol. Med.* 12, 2628-2643. doi: 10.1111/j.1582-4934.2008.00317.x
- [63] Nakamura, K., Ito, Y., Kawano, Y., Kurozumi, K., Kobune, M., Tsuda, H., et al. (2004). Antitumor effect of genetically engineered mesenchymal stem cells in a rat glioma model. *Gene Ther.* 11, 1155-1164. doi: 10.1038/sj.gt.3302276
- [64] NguyenThai, Q. A., Sharma, N., Luong do, H., Sodhi, S. S., Kim, J. H., Kim, N., et al. (2015). Targeted inhibition of osteosarcoma tumor growth by bone marrow-derived mesenchymal stem cells expressing cytosine deaminase/5-fluorocytosine in tumor-bearing mice. *J. Gene Med.* 17, 87-99. doi: 10.1002/jgm.2826

- [65] Nicolay, N. H., Lopez Perez, R., Ruhle, A., Trinh, T., Sisombath, S., Weber, K. J., et al. (2016). Mesenchymal stem cells maintain their defining stem cell characteristics after treatment with cisplatin. *Sci. Rep.* 6:20035. doi: 10.1038/srep20035
- [66] Nwabo Kamdje, A. H., Kamga, P. T., Simo, R. T., Vecchio, L., Seke Etet, P. F., Muller, J. M., et al. (2017). Mesenchymal stromal cells' role in tumor microenvironment: involvement of signaling pathways. *Cancer Biol. Med.* 14, 129-141. doi: 10.20892/j.issn.2095-3941.2016.0033
- [67] Otsu, K., Das, S., Houser, S. D., Quadri, S. K., Bhattacharya, S., and Bhattacharya, J. (2009). Concentration-dependent inhibition of angiogenesis by mesenchymal stem cells. *Blood* 113, 4197-4205. doi: 10.1182/blood-2008-09-176198
- [68] Pascucci, L., Cocce, V., Bonomi, A., Ami, D., Ceccarelli, P., Ciusani, E., et al. (2014). Paclitaxel is incorporated by mesenchymal stromal cells and released in exosomes that inhibit in vitro tumor growth: a new approach for drug delivery. *J. Control. Release* 192, 262-270. doi: 10.1016/j.jconrel.2014.07.042
- [69] Pessina, A., Cocce, V., Pascucci, L., Bonomi, A., Cavicchini, L., Sisto, F., et al. (2013). Mesenchymal stromal cells primed with Paclitaxel attract and kill leukaemia cells, inhibit angiogenesis and improve survival of leukaemia-bearing mice. *Br. J. Haematol.* 160, 766-778. doi: 10.1111/bjh.12196
- [70] Pessina, A., Piccirillo, M., Mineo, E., Catalani, P., Gribaldo, L., Marafante, E., et al. (1999). Role of SR-4987 stromal cells in the modulation of doxorubicin toxicity to in vitro granulocyte-macrophage progenitors (CFU-GM). *Life Sci.* 65, 513-523. doi: 10.1016/S0024-3205(99)00272-6
- [71] Pinilla, S., Alt, E., Abdul Khalek, F. J., Jotzu, C., Muehlberg, F., Beckmann, C., et al. (2009). Tissue resident stem cells produce CCL5 under the influence of cancer cells and thereby promote breast cancer cell invasion. *Cancer Lett.* 284, 80-85. doi: 10.1016/j.canlet.2009.04.013
- [72] Pokharel, D., Wijesinghe, P., Oenarto, V., Lu, J. F., Sampson, D. D., Kennedy, B. F., et al. (2016). Deciphering cell-to-cell communication in acquisition of cancer traits: extracellular membrane vesicles are regulators of tissue biomechanics. *OMICS* 20, 462-469. doi: 10.1089/omi.2016.0072
- [73] Ponte, A. L., Marais, E., Gallay, N., Langonne, A., Delorme, B., Herault, O., et al. (2007). The in vitro migration capacity of human bone marrow mesenchymal stem cells: comparison of chemokine and growth factor chemotactic activities. *Stem Cells* 25, 1737-1745. doi: 10.1634/stemcells.2007-0054
- [74] Qiao, L., Xu, Z., Zhao, T., Zhao, Z., Shi, M., Zhao, R. C., et al. (2008). Suppression of tumorigenesis by human mesenchymal stem cells in a hepatoma model. *Cell Res.* 18, 500-507. doi: 10.1038/cr.2008.40
- [75] Ramdasi, S., Sarang, S., and Viswanathan, C. (2015). Potential of mesenchymal stem cell based application in cancer. *Int. J. Hematol. Oncol. Stem Cell Res.* 9, 95-103.
- [76] Rattigan, Y., Hsu, J. M., Mishra, P. J., Glod, J., and Banerjee, D. (2010). Interleukin 6 mediated recruitment of mesenchymal stem cells to the hypoxic tumor milieu. *Exp. Cell Res.* 316, 3417-3424. doi: 10.1016/j.yexcr.2010.07.002
- [77] Ren, C., Kumar, S., Chanda, D., Chen, J., Mountz, J. D., and Ponnazhagan, S. (2008a). Therapeutic potential of mesenchymal stem cells producing interferon-alpha in a mouse melanoma lung metastasis model. *Stem Cells* 26, 2332-2338. doi: 10.1634/stemcells.2008-0084
- [78] Ren, C., Kumar, S., Chanda, D., Kallman, L., Chen, J., Mountz, J. D., et al. (2008b). Cancer gene therapy using mesenchymal stem cells expressing interferon-beta in a mouse prostate cancer lung metastasis model. *Gene Ther.* 15, 1446-1453. doi: 10.1038/gt.2008.101
- [79] Ringe, J., Strassburg, S., Neumann, K., Endres, M., Notter, M., Burmester, G. R., et al. (2007). Towards in situ tissue repair: human mesenchymal stem cells express chemokine receptors CXCR1, CXCR2 and CCR2, and migrate upon stimulation with CXCL8 but not CCL2. *J. Cell. Biochem.* 101, 135-146. doi: 10.1002/jcb.21172
- [80] Rivera-Cruz, C. M., Shearer, J. J., Figueiredo Neto, M., and Figueiredo, M. L. (2017). The immunomodulatory effects of mesenchymal stem cell polarization within the tumor microenvironment niche. *Stem Cells Int.* 2017:4015039. doi: 10.1155/2017/4015039
- [81] Rustad, K. C., and Gurtner, G. C. (2012). Mesenchymal stem cells home to sites of injury and inflammation. *Adv. Wound Care* 1, 147-152. doi: 10.1089/wound.2011.0314
- [82] Ryu, C. H., Park, S. H., Park, S. A., Kim, S. M., Lim, J. Y., Jeong, C. H., et al. (2011). Gene therapy of intracranial glioma using interleukin 12-secreting human umbilical cord blood-derived mesenchymal stem cells. *Hum. Gene Ther.* 22, 733-743. doi: 10.1089/hum.2010.187
- [83] Sage, E. K., Kolluri, K. K., McNulty, K., Lourenco Sda, S., Kalber, T. L., Ordidge, K. L., et al. (2014). Systemic but not topical TRAIL-expressing mesenchymal stem cells reduce tumour growth in malignant mesothelioma. *Thorax* 69, 638-647. doi: 10.1136/thoraxjnl-2013-204110
- [84] Sasaki, M., Abe, R., Fujita, Y., Ando, S., Inokuma, D., and Shimizu, H. (2008). Mesenchymal stem cells are recruited into wounded skin and contribute to wound repair by transdifferentiation into multiple skin cell type. *J. Immunol.* 180, 2581-2587. doi: 10.4049/jimmunol.180.4.2581
- [85] Schar, M. O., Diaz-Romero, J., Kohl, S., Zumstein, M. A., and Nesic, D. (2015). Platelet-rich concentrates differentially release growth factors and induce cell migration in vitro. *Clin. Orthop. Relat. Res.* 473, 1635-1643. doi: 10.1007/s11999-015-4192-2

- [86] Sharif, S., Ghahremani, M. H., and Soleimani, M. (2017). Delivery of exogenous miR-124 to glioblastoma multiform cells by Wharton's jelly mesenchymal stem cells decreases cell proliferation and migration, and confers chemosensitivity. *Stem Cell Rev.* doi: 10.1007/s12015-017-9788-3
- [87] Shi, S., Zhang, Q., Xia, Y., You, B., Shan, Y., Bao, L., et al. (2016). Mesenchymal stem cell-derived exosomes facilitate nasopharyngeal carcinoma progression. *Am. J. Cancer Res.* 6, 459-472.
- [88] Smith, C. L., Chaichana, K. L., Lee, Y. M., Lin, B., Stanko, K. M., O'Donnell, T., et al. (2015). Pre-exposure of human adipose mesenchymal stem cells to soluble factors enhances their homing to brain cancer. *Stem Cells Transl. Med.* 4, 239-251. doi: 10.5966/sctm.2014-0149
- [89] Son, B. R., Marquez-Curtis, L. A., Kucia, M., Wysoczynski, M., Turner, A. R., Ratajczak, J., et al. (2006). Migration of bone marrow and cord blood mesenchymal stem cells in vitro is regulated by stromal-derived factor-1-CXCR4 and hepatocyte growth factor-c-met axes and involves matrix metalloproteinases. *Stem Cells* 24, 1254-1264. doi: 10.1634/stemcells.2005-0271
- [90] Sordi, V., Malosio, M. L., Marchesi, F., Mercalli, A., Melzi, R., Giordano, T., et al. (2005). Bone marrow mesenchymal stem cells express a restricted set of functionally active chemokine receptors capable of promoting migration to pancreatic islets. *Blood* 106, 419-427. doi: 10.1182/blood-2004-09-3507
- [91] Srinivasula, S. M., Hegde, R., Saleh, A., Datta, P., Shiozaki, E., Chai, J., et al. (2001). A conserved XIAP-interaction motif in caspase-9 and Smac/DIABLO regulates caspase activity and apoptosis. *Nature* 410, 112-116. doi: 10.1038/35065125
- [92] Studeny, M., Marini, F. C., Champlin, R. E., Zompetta, C., Fidler, I. J., and Andreeff, M. (2002). Bone marrow-derived mesenchymal stem cells as vehicles for interferon-beta delivery into tumors. *Cancer Res.* 62, 3603-3608.
- [93] Su, H., Li, J., Osinska, H., Li, F., Robbins, J., Liu, J., et al. (2013). The COP9 signalosome is required for autophagy, proteasome-mediated proteolysis, and cardiomyocyte survival in adult mice. *Circ. Heart Fail.* 6, 1049-1057. doi: 10.1161/CIRCHEARTFAILURE.113.000338
- [94] Szegezdi, E., O'Reilly, A., Davy, Y., Vawda, R., Taylor, D. L., Murphy, M., et al. (2009). Stem cells are resistant to TRAIL receptor-mediated apoptosis. *J. Cell Mol. Med.* 13, 4409-4414. doi: 10.1111/j.1582-4934.2008.00522.x
- [95] Tsukamoto, S., Honoki, K., Fujii, H., Tohma, Y., Kido, A., Mori, T., et al. (2012). Mesenchymal stem cells promote tumor engraftment and metastatic colonization in rat osteosarcoma model. *Int. J. Oncol.* 40, 163-169. doi: 10.3892/ijo.2011.1220
- [96] Uchibori, R., Okada, T., Ito, T., Urabe, M., Mizukami, H., Kume, A., et al. (2009). Retroviral vector-producing mesenchymal stem cells for targeted suicide cancer gene therapy. *J. Gene Med.* 11, 373-381. doi: 10.1002/jgm.1313
- [97] Volarevic, V., Markovic, B. S., Gazdic, M., Volarevic, A., Jovicic, N., Arsenijevic, N., et al. (2018). Ethical and safety issues of stem cell-based therapy. *Int. J. Med. Sci.* 15, 36-45. doi: 10.7150/ijms.21666
- [98] Wang, G. X., Zhan, Y. A., Hu, H. L., Wang, Y., and Fu, B. (2012). Mesenchymal stem cells modified to express interferon-beta inhibit the growth of prostate cancer in a mouse model. *J. Int. Med. Res.* 40, 317-327. doi: 10.1177/147323001204000132
- [99] Wang, X. J., Xiang, B. Y., Ding, Y. H., Chen, L., Zou, H., Mou, X. Z., et al. (2017). Human menstrual blood-derived mesenchymal stem cells as a cellular vehicle for malignant glioma gene therapy. *Oncotarget* 8, 58309-58321. doi: 10.18632/oncotarget.17621
- [100] Wu, N., Zhang, Y. L., Wang, H. T., Li, D. W., Dai, H. J., Zhang, Q. Q., et al. (2016). Overexpression of hepatocyte nuclear factor 4alpha in human mesenchymal stem cells suppresses hepatocellular carcinoma development through Wnt/beta-catenin signaling pathway downregulation. *Cancer Biol. Ther.* 17, 558-565. doi: 10.1080/15384047.2016.1177675
- [101] Xia, L., Peng, R., Leng, W., Jia, R., Zeng, X., Yang, X., et al. (2015). TRAIL-expressing gingival-derived mesenchymal stem cells inhibit tumorigenesis of tongue squamous cell carcinoma. *J. Dent. Res.* 94, 219-228. doi: 10.1177/0022034514557815
- [102] Xie, C., Yang, Z., Suo, Y., Chen, Q., Wei, D., Weng, X., et al. (2017). Systemically infused mesenchymal stem cells show different homing profiles in healthy and tumor mouse models. *Stem Cells Transl. Med.* 6, 1120-1131. doi: 10.1002/sctm.16-0204
- [103] Xin, H., Kanehira, M., Mizuguchi, H., Hayakawa, T., Kikuchi, T., Nukiwa, T., et al. (2007). Targeted delivery of CX3CL1 to multiple lung tumors by mesenchymal stem cells. *Stem Cells* 25, 1618-1626. doi: 10.1634/stemcells.2006-0461
- [104] Xu, W. T., Bian, Z. Y., Fan, Q. M., Li, G., and Tang, T. T. (2009). Human mesenchymal stem cells (hMSCs) target osteosarcoma and promote its growth and pulmonary metastasis. *Cancer Lett.* 281, 32-41. doi: 10.1016/j.canlet.2009.02.022
- [105] Yan, C., Song, X., Yu, W., Wei, F., Li, H., Lv, M., et al. (2016). Human umbilical cord mesenchymal stem cells delivering sTRAIL home to lung cancer mediated by MCP-1/CCR2 axis and exhibit antitumor effects. *Tumour Biol.* 37, 8425-8435. doi: 10.1007/s13277-015-4746-7

- [106] Yan, Z., Zhuansun, Y., Chen, R., Li, J., and Ran, P. (2014a). Immunomodulation of mesenchymal stromal cells on regulatory T cells and its possible mechanism. *Exp. Cell Res.* 324, 65-74. doi: 10.1016/j.yexcr.2014.03.013
- [107] Yan, Z., Zhuansun, Y., Liu, G., Chen, R., Li, J., and Ran, P. (2014b). Mesenchymal stem cells suppress T cells by inducing apoptosis and through PD-1/B7-H1 interactions. *Immunol. Lett.* 162(1 Pt A), 248-255. doi: 10.1016/j.imlet.2014.09.013
- [108] Yang, L., Zhang, Y., Cheng, L., Yue, D., Ma, J., Zhao, D., et al. (2016). Mesenchymal stem cells engineered to secrete pigment epithelium-derived factor inhibit tumor metastasis and the formation of malignant ascites in a murine colorectal peritoneal carcinomatosis model. *Hum. Gene Ther.* 27, 267-277. doi: 10.1089/hum.2015.135
- [109] Yang, X., Du, J., Xu, X., Xu, C., and Song, W. (2014). IFN-gamma-secreting-mesenchymal stem cells exert an antitumor effect in vivo via the TRAIL pathway. *J. Immunol. Res.* 2014:318098. doi: 10.1155/2014/318098
- [110] Yang, Z. S., Tang, X. J., Guo, X. R., Zou, D. D., Sun, X. Y., Feng, J. B., et al. (2014). Cancer cell-oriented migration of mesenchymal stem cells engineered with an anticancer gene (PTEN): an imaging demonstration. *Onco Targets Ther.* 7, 441-446. doi: 10.2147/OTT.S59227
- [111] You, Q., Yao, Y., Zhang, Y., Fu, S., Du, M., and Zhang, G. (2015). Effect of targeted ovarian cancer therapy using amniotic fluid mesenchymal stem cells transfected with enhanced green fluorescent protein-human interleukin-2 in vivo. *Mol. Med. Rep.* 12, 4859-4866. doi: 10.3892/mmr.2015.4076
- [112] Yuan, Z., Kolluri, K. K., Gowers, K. H., and Janes, S. M. (2017). TRAIL delivery by MSC-derived extracellular vesicles is an effective anticancer therapy. *J. Extracell. Vesicles* 6:1265291. doi: 10.1080/20013078.2017.1265291
- [113] Yuan, Z., Kolluri, K. K., Sage, E. K., Gowers, K. H., and Janes, S. M. (2015). Mesenchymal stromal cell delivery of full-length tumor necrosis factor-related apoptosis-inducing ligand is superior to soluble type for cancer therapy. *Cytotherapy* 17, 885-896. doi: 10.1016/j.jcyt.2015.03.603
- [114] Zhang, B., Shan, H., Li, D., Li, Z. R., Zhu, K. S., and Jiang, Z. B. (2012). The inhibitory effect of MSCs expressing TRAIL as a cellular delivery vehicle in combination with cisplatin on hepatocellular carcinoma. *Cancer Biol. Ther.* 13, 1175-1184. doi: 10.4161/cbt.21347
- [115] Zhang, J., Hou, L., Wu, X., Zhao, D., Wang, Z., Hu, H., et al. (2016). Inhibitory effect of genetically engineered mesenchymal stem cells with Apoptin on hepatoma cells in vitro and in vivo. *Mol. Cell. Biochem.* 416, 193-203. doi: 10.1007/s11010-016-2707-0
- [116] Zhao, W. H., Cheng, J. X., Shi, P. F., and Huang, J. Y. (2011). Human umbilical cord mesenchymal stem cells with adenovirus-mediated interleukin 12 gene transduction inhibits the growth of ovarian carcinoma cells both in vitro and in vivo. *Nan Fang Yi Ke Da Xue Xue Bao* 31, 903-907.
- [117] Zhu, Y., Cheng, M., Yang, Z., Zeng, C. Y., Chen, J., Xie, Y., et al. (2014). Mesenchymal stem cell-based NK4 gene therapy in nude mice bearing gastric cancer xenografts. *Drug Des. Dev. Ther.* 8, 2449-2462. doi: 10.2147/DDDT.S71466